

The locus of real multiplication and the Schottky locus

Matt Bainbridge and Martin Möller

December 14, 2011

Contents

1	Introduction	1
2	Boundary of the real multiplication locus	3
3	Properties of the RM-tori	9
4	Reduction to the boundary condition	12
5	Cross-ratios for nice boundary strata without separating nodes	15
6	Checking nice boundary strata without separating nodes	17
7	Strata with separating curves	20

1 Introduction

Are there locally symmetric subvarieties in the moduli space of abelian varieties, whose generic point lies in the image of the moduli space of curves, i.e. in the Schottky locus? This question was raised by Oort, motivated by the Conjecture of Coleman that for g large enough there are only finitely many curves of genus g with complex multiplication.

The first important contributions to this problem were made by Hain [Hai99], whose results were subsequently refined by de Jong and Zhang [dJZ07], see also [MO] for a survey. In this paper we do not consider general locally symmetric subvarieties (or Shimura subvarieties) but restrict to the case of Hilbert modular varieties. They parametrize abelian varieties with real multiplication. So our main result is a statement about components of the real multiplication locus.

Theorem 1.1. *There is no component of the real multiplication locus in the moduli space of four-dimensional abelian varieties \mathcal{A}_4 that lies generically in the image of the moduli space of curves \mathcal{M}_4 .*

In [dJZ07] the analogous theorem is proved for genus greater than four with part of the genus four case still left open. Together, the two results imply:

Corollary 1.2. *For every genus g and every component of the real multiplication locus in the moduli space of g -dimensional abelian varieties \mathcal{A}_g , the generic*

point of the component does not lie in the image of the moduli space of curves \mathcal{M}_g .

Besides completing the work of [dJZ07], we believe our method of proof is interesting for the following reason.

The proofs in [dJZ07] and [Hai99] ultimately rely on a theorem of Farb and Masur [FM98] that mapping class groups do not contain fundamental groups of lattices in higher rank Lie groups. Consequently, if the generic point of a component of the real multiplication locus lies in the moduli space of curves, then the Torelli map must modify the fundamental group of the real multiplication locus, either by ramification along the hyperelliptic locus, or by the locus of decomposable abelian varieties, which is disjoint from the Schottky locus. Whenever this can be ruled out, e.g. by showing that the codimension of this intersection is at least two, one has the desired contradiction.

The proof here, on the contrary, relies on a study at the boundary of the moduli space of curves. Since Hilbert modular varieties are not compact, we may study their closure in the Deligne-Mumford compactification. A counterexample to the theorem in genus four must have a component of dimension three in some Deligne-Mumford boundary stratum of the moduli space of curves.

These closures of Hilbert modular varieties were analyzed in [BM]. If one uses cross-ratios as degenerate period coordinates, the closure of a Hilbert modular variety is contained in a subtorus of an ambient algebraic torus which we call the *RM-torus*. So a first try to rule out counterexamples to the main theorem is to check if the images of Deligne-Mumford boundary strata in cross-ratio coordinates contain tori of sufficiently large dimension.

In fact, they do contain such large tori, but only when the tori are very degenerate, e.g. lying completely in a coordinate hyperplane. The heart of the paper consists in showing that RM-tori do not have this property. For that purpose, following the ideas in [BM], we provide the (dual graph of the) boundary stratum with *weights* given by the residues of an eigenform for real multiplication. Only if the weights satisfy the restrictive condition of being admissible, can the boundary stratum lie in the closure of the Hilbert modular variety. This condition is recalled in Theorem 2.1.

The obvious refinement of the above theorem, to understand the *dimension* of the intersection of the real multiplication locus in \mathcal{A}_g with \mathcal{M}_g (say for large g) is still an open problem. The techniques in this paper could conceivably contribute to the solution of this problem, which is not tractable by methods based on the lattice properties of fundamental group.

Section 3 derives the key properties of RM-tori, e.g. a method to calculate their intersection dimension with subtori in terms of the weights. Section 4 contains the details of the strategy outlined above. The main theorem is reduced in that section to showing for a list of relevant Deligne-Mumford boundary strata that the RM-tori are not contained in the image of the stratum under the cross-ratio maps. In Section 5 we show that cross-ratios are indeed coordinates near the boundary of a relevant boundary stratum if this boundary stratum does not parametrize curves with a separating node. In Section 6 we give graph-theoretic

criteria for the desired non-containment statement and thus complete the argument for relevant boundary strata parameterizing curves without a separating node. In the last section we deal with boundary strata that parametrize curves with a separating node and reduce this to a case previously dealt with.

2 Boundary of the real multiplication locus

In this section, we summarize properties of the real multiplication locus and their boundaries which will be needed in later sections. Throughout this section, F will denote a totally real number field of degree g , and \mathcal{O} will denote an order in F (a subring which has rank g as an abelian group). We start with some motivation and remarks explaining the approach used in [BM] to compute the boundary of the real multiplication locus. The goal of this section is to provide the notation and background to state this result as Theorem 2.1 below.

Our approach to studying boundaries of Hilbert modular varieties differs from the standard approach of toroidal compactifications. For those more familiar with toroidal compactifications, we compare briefly these two approaches at the end of this section.

The real multiplication locus. We denote by $\mathcal{RA}_{\mathcal{O}} \subset \mathcal{A}_g$ the locus of abelian varieties which have real multiplication by \mathcal{O} . This locus is an immersed quotient of a Hilbert modular variety by a finite group of automorphisms. We denote by $\mathcal{RM}_{\mathcal{O}} \subset \mathcal{M}_g$ the locus of Riemann surfaces whose Jacobians have real multiplication by \mathcal{O} . In other words, $\mathcal{RM}_{\mathcal{O}} = t^{-1}(\mathcal{RA}_{\mathcal{O}})$, where $t: \mathcal{M}_g \rightarrow \mathcal{A}_g$ is the Torelli map.

Irreducible components. Given an abelian variety A with real multiplication \mathcal{O} , the homology group $H_1(A; \mathbb{Z})$ has the structure of an \mathcal{O} -module with a compatible symplectic structure, and the isomorphism classes of such modules parametrize the irreducible components of $\mathcal{RA}_{\mathcal{O}}$.

More precisely, consider a torsion-free \mathcal{O} -module M . The *rank* of M is the dimension of $M \otimes \mathbb{Q}$ as a vector space over F . We say that M is *proper* if the \mathcal{O} -module structure doesn't extend to a larger order. A *symplectic \mathcal{O} -module* is a torsion-free \mathcal{O} -module equipped with a unimodular symplectic form satisfying $\langle x, \lambda y \rangle = \langle \lambda x, y \rangle$ for each $x, y \in M$ and $\lambda \in \mathcal{O}$.

Given a torsion-free, proper, rank-two, symplectic \mathcal{O} -module M , we define $\mathcal{RA}_{\mathcal{O}, M} \subset \mathcal{RA}_{\mathcal{O}}$ to be the locus of abelian varieties whose first homology is isomorphic to M as a symplectic \mathcal{O} -module. $\mathcal{RA}_{\mathcal{O}, M}$ is isomorphic to a finite quotient of a Hilbert modular variety \mathbb{H}^g/Γ for some Γ commensurable with $\mathrm{SL}_2(\mathcal{O})$, so it is irreducible. Thus the irreducible components of $\mathcal{RA}_{\mathcal{O}}$ are parametrized by isomorphism classes of such M .

We will use the notation $\mathcal{RM}_{\mathcal{O}, M} = t^{-1}(\mathcal{RA}_{\mathcal{O}, M}) \subset \mathcal{M}_g$.

Cusps. A lattice \mathcal{I} in F is a rank g additive subgroup $\mathcal{I} \subset F$. The *coefficient ring* of \mathcal{I} is the order $\mathcal{O}_{\mathcal{I}}$ defined by

$$\mathcal{O}_{\mathcal{I}} = \{a \in F : ax \in \mathcal{I} \text{ for all } x \in \mathcal{I}\}.$$

The *inverse different* of \mathcal{I} is the lattice \mathcal{I}^{\vee} defined by

$$\mathcal{I}^{\vee} = \{x \in F : \langle x, y \rangle \in \mathbb{Z} \text{ for all } y \in \mathcal{I}\},$$

also having coefficient ring $\mathcal{O}_{\mathcal{I}}$. Here and throughout the paper we use the notation $\langle x, y \rangle = \text{Tr}(xy)$ for the trace pairing.

Consider a rank-two symplectic \mathcal{O} -module M and a lattice \mathcal{I} whose coefficient ring contains \mathcal{O} . An exact sequence of \mathcal{O} -modules,

$$0 \rightarrow \mathcal{I} \rightarrow M \rightarrow \mathcal{I}^{\vee} \rightarrow 0, \tag{2.1}$$

expresses M as an extension of \mathcal{I}^{\vee} by \mathcal{I} . The sequence (2.1) splits as a sequence of abelian groups, yielding a group isomorphism $\mathcal{I} \oplus \mathcal{I}^{\vee} \rightarrow M$. The module $\mathcal{I} \oplus \mathcal{I}^{\vee}$ carries a natural symplectic structure, defined by

$$\langle (a, b), (c, d) \rangle = \text{Tr}(ad - bc).$$

This induces a symplectic structure on M which does not depend on the choice of splitting of (2.1). We define $E(\mathcal{I}, M)$ to be the set of isomorphism classes of extensions (2.1) such that the induced symplectic form on M agrees with the given one.

Stable forms. Consider a stable curve X , and let $X' \subset X$ be the complement of the nodes. A *stable form* on X is a holomorphic one-form on X' which has at worst simple poles at the cusps of X' , with opposite residues at two cusps which share a node.

Weighted stable curves. Consider a lattice \mathcal{I} in a totally real number field F of degree g . An \mathcal{I} -weighted stable curve is an arithmetic genus g , geometric genus 0 stable curve X , together with an element of \mathcal{I} assigned to each cusp of X' (the complement of the nodes), called the weight of that cusp, subject to the following restrictions:

- Cusps of X' sharing a node have opposite weight.
- The sum of weights of a component of X' is zero.
- The weights span \mathcal{I} .

One could think of a weighted stable curve as a curve together with a stable form whose residues belong to F .

When we do not care to specify the lattice \mathcal{I} , we may speak of a F -weighted stable curve, or just a weighted stable curve.

Two weighted stable curves are *isomorphic* (resp. *topologically equivalent*) if there is a weight preserving isomorphism (resp. homeomorphism) between the underlying stable curves.

If we don't want to specify the ideal (or the field) the weights span we just talk of *F-weighted* (or just weighted) stable curves, with the implicit meaning that the weights span a lattice \mathcal{I} .

Weighted boundary strata. We define an \mathcal{I} -weighted boundary stratum \mathcal{S} to be the moduli space of \mathcal{I} -weighted stable curves which are topologically equivalent to some fixed weighted stable curve X . If X has m components, each having n_i cusps with all weights distinct, then the stratum \mathcal{S} is isomorphic to

$$\prod_i \mathcal{M}_{0,n_i},$$

where $\mathcal{M}_{0,n}$ is the moduli space of n labeled points on \mathbb{P}^1 . If some weights coincide, the stratum may be a quotient of this product. There is a canonical morphism $\mathcal{S} \rightarrow \overline{\mathcal{M}}_g$ which forgets the weights.

An \mathcal{I} -weighted boundary stratum, or equivalently the topological type of a weighted stable curve may be encoded by a directed graph with edges weighted by elements of \mathcal{I} . Given a stratum \mathcal{S} parameterizing weighted curves topologically equivalent to X , we write $\Gamma(\mathcal{S})$ for the graph with one vertex for each component of X , with an edge joining two vertices if the corresponding components are joined by a node (the *dual graph*). Contrary to usual practice, we allow graphs where an edge joins a vertex to itself, or where multiple edges join the same pair of vertices. We label each edge with the weight of the corresponding node and an arrow pointing to the component with that weight (as opposed to its negative). We call such an object an \mathcal{I} -weighted graph. Two graphs which are related by changing the orientation of an edge and simultaneously the sign of its weight represent the same weighted boundary stratum, and we regard two such weighted graphs to be the same.

A *degeneration* of a weighted boundary stratum \mathcal{S} is a stratum obtained by pinching one or more simple closed curves on stable curves parametrized by \mathcal{S} . A degeneration \mathcal{S}' of \mathcal{S} can be regarded as part of the boundary of the Deligne-Mumford compactification of \mathcal{S} . On the level of dual graphs, degenerations of \mathcal{S} are obtained by gluing an edge into a vertex v of $\Gamma(\mathcal{S})$. More precisely, we replace the vertex v two vertices v_1 and v_2 joined by an edge e , with each edge meeting v now meeting either v_1 or v_2 . We assign e the unique weight which is consistent with the axioms of a weighted graph.

Periods. Consider an \mathcal{I} -weighted boundary stratum \mathcal{S} . We recall here a coordinate-free analogue of classical period matrices for weighted stable curves introduced in [BM].

Given any ring R and module M over R , we define $\text{Sym}_R(M)$ to be the submodule of $M \otimes_R M$ fixed by the involution $\theta(x \otimes y) = y \otimes x$. We define

$\mathbf{S}_R(M)$ to be the quotient of $M \otimes_R M$ by the submodule generated by the relations $\theta(z) - z$.

We identify the field F with its dual via the trace pairing; thus the vector spaces $\text{Sym}_{\mathbb{Q}}(F)$ and $\mathbf{S}_{\mathbb{Q}}(F)$ are dual via the pairing

$$\langle a \otimes b, c \otimes d \rangle = \langle a, c \rangle \langle b, d \rangle,$$

as are the groups $\text{Sym}_{\mathbb{Z}}(\mathcal{I})$ and $\mathbf{S}_{\mathbb{Z}}(\mathcal{I}^{\vee})$.

Let $W(\mathcal{S}) \subset \text{Sym}_{\mathbb{Q}}(F)$ be the subspace generated by the elements $r \otimes r$ for r running over the weights of \mathcal{S} . Let $N(\mathcal{S}) \subset \mathbf{S}_{\mathbb{Q}}(F)$ be the annihilator of $W(\mathcal{S})$.

We defined in [BM] a homomorphism

$$\Psi: N(\mathcal{S}) \cap \mathbf{S}_{\mathbb{Z}}(\mathcal{I}^{\vee}) \rightarrow \text{Hol}^*(\mathcal{S}),$$

where the image is the multiplicative group of nonzero holomorphic functions on \mathcal{S} . Each $\Psi(x)$ is a holomorphic function on \mathcal{S} which arises as a limit of an exponential of a classical period matrix entry. We describe here $\Psi(x)$ when x is an elementary tensor $\alpha \otimes \beta$, and refer the reader to [BM] for a more careful definition.

Consider $\alpha \otimes \beta \in N(\mathcal{S}) \cap \mathbf{S}_{\mathbb{Z}}(\mathcal{I}^{\vee})$ and a weighted stable curve $X \in \mathcal{S}$. Pairing α with the weights of X associates an integer to each cusp of X . There is a unique stable form on X with these residues at the cusps, which we call ω_X . Similarly, pairing β with the weights associates an integer to each cusp, and we may choose a cycle γ on X whose algebraic intersection number with each node is given by these integers. We define

$$\Psi(\alpha \otimes \beta)(X) = e^{\int_{\gamma} \omega_X}. \quad (2.2)$$

Since $\alpha \otimes \beta$ belongs to $N(\mathcal{S})$, we may choose γ to not pass through any nodes at which ω_X has a pole, so this integral is finite and well-defined. It may be checked directly that $\Psi(\alpha \otimes \beta) = \Psi(\beta \otimes \alpha)$, or more conceptually this follows from the symmetry of period matrices of nonsingular curves by a degeneration argument.

The function $\Psi(x)$ is always a product of various cross-ratios of points on components the stable curve (see Lemma 2.2).

The necessary condition. We now recall the necessary condition for a stable curve to lie in the boundary of the real multiplication locus $\mathcal{RM}_{\mathcal{O}}$.

Consider an \mathcal{I} -weighted boundary stratum \mathcal{S} . In $\mathbf{S}_{\mathbb{Q}}(F)$, we define the cone

$$C(\mathcal{S}) = \{x \in \mathbf{S}_{\mathbb{Q}}(F) : \langle x, r \otimes r \rangle \geq 0 \text{ for each weight } r \text{ of } \mathcal{S}\}.$$

The space $F \otimes_{\mathbb{Q}} F$ has the structure of an F -bimodule. We define

$$\Lambda^1 = \{x \in F \otimes_{\mathbb{Q}} F : \lambda \cdot x = x \cdot \lambda \text{ for each } \lambda \in F\}.$$

In fact, Λ^1 is contained in $\text{Sym}_{\mathbb{Q}}(F) \subset F \otimes_{\mathbb{Q}} F$ (see [BM, Proposition 5.1]). We define $\text{Ann}(\Lambda^1) \subset \mathbf{S}_{\mathbb{Q}}(F)$ to be the annihilator of Λ^1 .

We say that the weighted stratum \mathcal{S} is *admissible* if

$$C(\mathcal{S}) \cap \text{Ann}(\Lambda^1) \subset N(\mathcal{S}).$$

We associate to an admissible stratum \mathcal{S} various algebraic tori. We define the *ambient torus* $A_{\mathcal{S}}$ by

$$A_{\mathcal{S}} = \text{Hom}_{\mathbb{Z}}(N(\mathcal{S}) \cap \mathbf{S}_{\mathbb{Z}}(\mathcal{I}^{\vee}), \mathbb{G}_m).$$

(Readers unfamiliar with algebraic groups should regard \mathbb{G}_m as the multiplicative group of nonzero complex numbers.) The homomorphism $\Psi: N(\mathcal{S}) \cap \mathbf{S}_{\mathbb{Z}}(\mathcal{I}^{\vee}) \rightarrow \text{Hol}^*(\mathcal{S})$ determines a dual morphism $\text{CR}: \mathcal{S} \rightarrow A_{\mathcal{S}}$.

There is a surjective map of algebraic tori:

$$p: A_{\mathcal{S}} \rightarrow \text{Hom}(N(\mathcal{S}) \cap \text{Ann}(\Lambda^1) \cap \mathbf{S}_{\mathbb{Z}}(\mathcal{I}^{\vee}), \mathbb{G}_m). \quad (2.3)$$

We define the *real multiplication torus* (or *RM-torus*) $T_{\mathcal{S}}$ to be the subtorus $T_{\mathcal{S}} = p^{-1}(0) \subset A_{\mathcal{S}}$. More generally, there is a function

$$q: E(\mathcal{I}, M) \rightarrow \text{Hom}(N(\mathcal{S}) \cap \text{Ann}(\Lambda^1) \cap \mathbf{S}_{\mathbb{Z}}(\mathcal{I}^{\vee}), \mathbb{G}_m),$$

with image in the set of torsion points, which is defined in [BM, §5]; we refer the reader to that paper for the definition, as it is not needed here. Given an extension $E \in E(\mathcal{I}, M)$, we define the *translated RM-torus* $T_{\mathcal{S}, E}$ by $T_{\mathcal{S}, E} = p^{-1}(q(E)) \subset A_{\mathcal{S}}$. Given an extension E , we define the subvariety $\mathcal{RS}_E \subset \mathcal{S}$ to be the inverse image of $T_{\mathcal{S}, E}$ under CR .

We can now state our necessary condition for a geometric genus zero stable curve to lie in the boundary of the real multiplication locus. See [BM, §5] for the proof.

Theorem 2.1. *If a geometric genus zero stable curve $X \in \overline{\mathcal{M}}_g$ lies in the boundary of $\mathcal{RM}_{\mathcal{O}, M}$, then there is a lattice $\mathcal{I} \subset F$ whose coefficient ring contains \mathcal{O} and an extension $E \in E(\mathcal{I}, M)$ such that X is in the image of \mathcal{RS}_E under the forgetful map $\mathcal{S} \rightarrow \overline{\mathcal{M}}_g$ for some admissible \mathcal{I} -weighted boundary stratum \mathcal{S} .*

The idea of the proof of this theorem is quite simple. Each $a \in \mathbf{S}_{\mathbb{Z}}(\mathcal{I}^{\vee})$ determines a meromorphic function $\Psi(a)$ on a neighborhood of \mathcal{S} (or a suitable cover of \mathcal{M}_g) which is a product of exponentials of period matrix entries. This $\Psi(a)$ is nonzero and finite on \mathcal{S} exactly when $a \in N(\mathcal{S})$, and it is constant on $\mathcal{RM}_{\mathcal{O}, M}$ exactly when $a \in \text{Ann}(\Lambda^1)$. These $\Psi(a)$ restricted to \mathcal{S} cut out the subvariety \mathcal{RS}_E . Finally, if the weights are not admissible, then one can find $\Psi(a)$ which is a nonzero constant on $\mathcal{RM}_{\mathcal{O}, M}$, but which vanishes on \mathcal{S} , hence these subvarieties must be disjoint.

Loops in weighted graphs. Consider a geometric genus zero weighted stable curve X lying in a stratum \mathcal{S} . There is a natural bijection between loops in the weighted graph $\Gamma(\mathcal{S})$ and homotopy classes of loops on X . We now describe a method to construct elements of $N(\mathcal{S})$ from pairs of loops in the weighted graph $\Gamma(\mathcal{S})$.

We say that two loops in a graph are *edge-disjoint*, if they do not share an edge. If they do not share a vertex, we call them *vertex-disjoint*. We say that a loop is *simple* if it meets each vertex at most once.

To a loop γ in $\Gamma(\mathcal{S})$, define a functional $\lambda^*(\gamma) \in \text{Hom}_{\mathbb{Q}}(F, \mathbb{Q})$ so that for any edge e having weight r , $\lambda^*(\gamma)(r)$ is the number of times γ traverses e in the positive direction minus the number of times γ traverses e in the negative direction. This defines a functional on F as the weights span F . It is linear and well defined by the properties of a weighted stable curve.

We define $\lambda(\gamma) \in F$ to be the unique element such that

$$\langle \lambda(\gamma), x \rangle = \lambda^*(\gamma)(x)$$

for all $x \in F$. If we think of γ as a loop on the stable curve X , and if n is a node of X having weight r , then $\langle \lambda(\gamma), r \rangle$ is the algebraic intersection number of γ with n . We may also regard λ as an isomorphism $\lambda: H_1(\Gamma(\mathcal{S}); \mathbb{Z}) \rightarrow \mathcal{I}^{\vee}$.

We can calculate $\lambda(\gamma)$ explicitly as follows. Choose edges e_1, \dots, e_g of $\Gamma(\mathcal{S})$ having corresponding weights $r_1, \dots, r_g \in F$ such that these r_i form a basis of F over \mathbb{Q} . Let s_1, \dots, s_g be the dual basis of F with respect to the trace pairing. Then $\lambda(\gamma) = \sum n_i s_i$, where n_i is the number of times γ traverses e_i , crossings with the opposite orientation counted negatively.

If a simple loop γ passes through a vertex v , we denote by $\gamma^{\text{in}}(v)$ (resp. $\gamma^{\text{out}}(v)$) the incoming (resp. outgoing) marked point on the component of the stable curve corresponding to v .

Given $a, b, c, d \in \mathbb{C}$, recall that their cross-ratio is defined by

$$[a, b, c, d] = \frac{(a-c)(b-d)}{(a-d)(b-c)}.$$

Lemma 2.2. *Suppose the loops γ_1 and γ_2 in $\Gamma(\mathcal{S})$ are edge-disjoint. Then the corresponding element $a = \lambda(\gamma_1) \otimes \lambda(\gamma_2) \in \mathbf{S}(F)$ lies in $N(\mathcal{S})$. Moreover, if the γ_i are simple, then the function $\Psi(a)$ is a product of cross-ratios*

$$\Psi(a) = \prod_{v \in \gamma_1 \cap \gamma_2} [\gamma_1^{\text{out}}(v), \gamma_1^{\text{in}}(v), \gamma_2^{\text{out}}(v), \gamma_2^{\text{in}}(v)].$$

Proof. Consider an edge e having weight r . Since the loops share no edges, one of the γ_i does not pass through e . We then have $\langle \lambda(\gamma_i), r \rangle = 0$, so

$$\langle \lambda(\gamma_1) \otimes \lambda(\gamma_2), r \otimes r \rangle = 0.$$

It follows that $\lambda(\gamma_1) \otimes \lambda(\gamma_2) \in N(\mathcal{S})$.

We consider now γ_1 and γ_2 as loops on a stable curve X in \mathcal{S} . There is a unique stable form ω on X with poles of residue -1 at each $\gamma_1^{\text{out}}(v)$ and residue 1 at each $\gamma_1^{\text{in}}(v)$. By definition, $\Psi(a) = e^{\int_{\gamma_2} \omega}$.

Fix a component of X corresponding to a vertex v , which we identify with the Riemann sphere punctured at finitely many points. Normalizing so that

$\gamma_1^{\text{in}}(v) = 0$ and $\gamma_2^{\text{out}}(v) = \infty$, we have $\omega = \frac{dz}{z}$ on this component and the corresponding term of $\int_{\gamma_2} \omega$ is

$$\int_{\gamma_2^{\text{out}}(v)}^{\gamma_2^{\text{in}}(v)} \frac{dz}{z} = \log \frac{\gamma_2^{\text{in}}(v)}{\gamma_2^{\text{out}}(v)} = \log[\gamma_1^{\text{out}}(v), \gamma_1^{\text{in}}(v), \gamma_2^{\text{out}}(v), \gamma_2^{\text{in}}(v)]. \quad \blacksquare$$

The boundary and toroidal compactifications. The moduli space of abelian varieties \mathcal{A}_g admits many toroidal compactifications depending on an admissible fan decomposition of the space of semi-positive-definite quadratic forms on \mathbb{Z}^g . Among the many toroidal compactifications, the second Voronoi decomposition $\overline{\mathcal{A}}_g^{\text{Vor}}$ is very appealing in this context, since by the work of Mumford and Namikawa the Torelli map extends to a map $\bar{t}: \overline{\mathcal{M}}_g \rightarrow \overline{\mathcal{A}}_g^{\text{Vor}}$ (see [Nam76] and e.g. [Ale04] for a recent exposition on these topics).

We will deduce the main theorem from properties of the intersection of $\overline{\mathcal{RM}}_{\mathcal{O}}$ with the boundary of $\overline{\mathcal{M}}_g$ developed in [BM] which we summarized above. An alternative approach would have been to describe the closure of $\mathcal{RA}_{\mathcal{O}}$ inside $\overline{\mathcal{A}}_g^{\text{Vor}}$ and then to take the intersection with $\bar{t}(\overline{\mathcal{M}}_g)$, taking special care on when \bar{t} is not an embedding. This procedure cannot circumvent the total complexity of the problem. The reader may want to look at the boundary of the real multiplication locus of smallest discriminant in genus 3, shown as Figure 3 in [BM]. Already in this simplest case, the structure of the boundary is rather complicated.

On the other hand, the possibility of computing in a toroidal compactification makes the structure of the result very plausible. If a stable curve has no separating nodes, cross ratios will define an embedding of a corresponding stratum \mathcal{S} of $\partial\overline{\mathcal{M}}_g$ into the ambient torus $A_{\mathcal{S}}$. In the period coordinates near the boundary of a toroidal compactification, real multiplication is a linear subvariety, and consequently $\overline{\mathcal{RM}}_{\mathcal{O}}$ is cut out by the translate of a subtorus, called the real multiplication torus $T_{\mathcal{S}}$ above.

An explicit toric description of the compactification the real multiplication locus in the genus two defined in [Bai07] in terms of a fan is provided by [MZ11].

3 Properties of the RM-tori

We now study in more detail the tori $T_{\mathcal{S}}$ and $A_{\mathcal{S}}$ introduced in §2, computing their dimension, as well of the dimension of the intersection of $T_{\mathcal{S}}$ with various subtori of $A_{\mathcal{S}}$.

Theorem 3.1. *Consider a F -weighted boundary stratum \mathcal{S} having genus g , among whose weights are exactly n distinct weights (up to sign) r_1, \dots, r_n . Then the elements $r_1 \otimes r_1, \dots, r_n \otimes r_n$ of $\text{Sym}_{\mathbb{Q}}(F)$ are linearly independent over \mathbb{Q} .*

Equivalently, $N(\mathcal{S})$ and $A_{\mathcal{S}}$ have dimension $g(g+1)/2 - n$.

Proof. The equivalence of these statements is clear from the definition of $N(\mathcal{S})$ and $A_{\mathcal{S}}$. Suppose first that \mathcal{S} is an irreducible stratum (parameterizing irreducible stable curves). There are g distinct weights r_1, \dots, r_g which form a

basis of F over \mathbb{Q} , and the $r_i \otimes r_i$ are then linearly independent in $\text{Sym}_{\mathbb{Q}}(F)$ (as is true for any basis of a vector space).

We now show that if the claim holds for some weighted stratum \mathcal{S} , then it holds for any degeneration \mathcal{S}' of \mathcal{S} obtained by pinching a single curve. The claim then follows for all strata by induction.

If the new node of \mathcal{S}' has the same weight as some node of \mathcal{S} , then $N(\mathcal{S}) = N(\mathcal{S}')$, and we are done. Now suppose the weight of the new node is distinct. To finish the proof, we must find an element of $N(\mathcal{S})$ which does not belong to $N(\mathcal{S}')$. In the weighted graph $\Gamma(\mathcal{S}')$, let e be the edge corresponding to the new node. Using the interpretation of pairs of loops on $\Gamma(\mathcal{S}')$ as elements of $\mathbf{S}_{\mathbb{Q}}(F)$ from §2, it suffices to find a pair of loops on $\Gamma(\mathcal{S}')$ which both contain the edge e and have no other edges in common.

Let G be the graph obtained by deleting e from $\Gamma(\mathcal{S}')$, and let p and q be the distinct vertices of G which were joined by e . Since the weight of e is distinct from the weights of the other edges of $\Gamma(\mathcal{S}')$, there is no edge of $\Gamma(\mathcal{S}')$, which jointly with e separates $\Gamma(\mathcal{S}')$. Thus G is not separated by any of its edges. It then follows from Menger's theorem (see [BM76]) that there are two edge-disjoint paths on G joining p to q . These paths yield the required pair of loops on $\Gamma(\mathcal{S}')$. ■

Corollary 3.2. *For any F -weighted boundary stratum \mathcal{S} , the cone $C(\mathcal{S}) \subset \mathbf{S}_{\mathbb{Q}}(F)$ strictly contains the subspace $N(\mathcal{S})$.*

Proof. Let r_i be the weights of \mathcal{S} . As the $r_i \otimes r_i$ are linearly independent in $\text{Sym}_{\mathbb{Q}}(F)$, we may find some $t \in \mathbf{S}_{\mathbb{Q}}(F)$ which pairs positively with $r_1 \otimes r_1$ and trivially with the other $r_i \otimes r_i$. This t lies in $C(\mathcal{S})$ but not $N(\mathcal{S})$. ■

We now turn to the dimension of the RM-torus $T_{\mathcal{S}}$.

Lemma 3.3. *The subspace $\Lambda^1 \subset \text{Sym}_{\mathbb{Q}}(F)$ has dimension g .*

Proof. Under the identification of $F \otimes_{\mathbb{Q}} F$ with $\text{Hom}_{\mathbb{Q}}(F, F)$ induced by the trace pairing, Λ^1 corresponds to $\text{Hom}_F(F, F)$. ■

Proposition 3.4. *For any admissible F -weighted boundary stratum \mathcal{S} of genus g , the RM-torus $T_{\mathcal{S}}$ has dimension at most $g - 1$.*

Proof. From the definition of $T_{\mathcal{S}}$, we have

$$\dim T_{\mathcal{S}} = \dim N(\mathcal{S}) - \dim(N(\mathcal{S}) \cap \text{Ann}(\Lambda^1)).$$

Under the quotient map $\mathbf{S}_{\mathbb{Q}}(F) \rightarrow \mathbf{S}_{\mathbb{Q}}(F)/N(\mathcal{S})$, the images of $\text{Ann}(\Lambda^1)$ and $C(\mathcal{S})$ have trivial intersection by the admissibility of \mathcal{S} . By Corollary 3.2, the image of $C(\mathcal{S})$ is nontrivial. It follows that the image of $\text{Ann}(\Lambda^1)$ is not all of $\mathbf{S}_{\mathbb{Q}}(F)/N(\mathcal{S})$. Equivalently,

$$\dim \text{Ann}(\Lambda^1) - \dim(\text{Ann}(\Lambda^1) \cap N(\mathcal{S})) < \dim \mathbf{S}_{\mathbb{Q}}(F) - \dim N(\mathcal{S}).$$

As $\text{Ann}(\Lambda^1)$ has codimension g in $\mathbf{S}_{\mathbb{Q}}(F)$ by Lemma 3.3, the desired inequality follows. ■

There are examples where $T_{\mathcal{S}}$ has dimension less than $g - 1$; see [BM, Appendix A].

Choose a basis r_1, \dots, r_g of F , and let s_1, \dots, s_g be the dual basis with respect to the trace pairing. We define

$$\epsilon = \sum_{i=1}^g r_i \otimes s_i \in F \otimes_{\mathbb{Q}} F.$$

Proposition 3.5. *The element ϵ lies in Λ^1 and does not depend on the choice of the basis of F . Thus we have $\Lambda^1 = \{x\epsilon : x \in F\}$. For every $x \in F$ and $s \otimes t \in \mathbf{S}_{\mathbb{Q}}(F)$, we have the pairing*

$$\langle x\epsilon, s \otimes t \rangle = \mathrm{Tr}_{\mathbb{Q}}^F(xst) \quad (3.1)$$

Proof. See [BM, Lemma 6.2]. This lemma only calculates the pairing $\langle x\epsilon, t \otimes t \rangle$, but the proof of the more general statement is identical. ■

We define the evaluation map $\mathrm{ev}: \mathbf{S}_{\mathbb{Q}}(F) \rightarrow F$ by $\mathrm{ev}(s \otimes t) = st$.

Corollary 3.6. *The annihilator $\mathrm{Ann}(\Lambda^1)$ is the kernel of ev .*

Proof. If $\alpha \in \mathrm{Ann}(\Lambda^1)$, then we have by Proposition 3.5 that

$$0 = \langle x\epsilon, \alpha \rangle = \mathrm{Tr}(\mathrm{ev}(\alpha)x)$$

for all $x \in F$. Since the trace pairing is nondegenerate, it follows that $\mathrm{ev}(\alpha) = 0$. ■

By the definition of $A_{\mathcal{S}}$, we have the identification $\chi(A_{\mathcal{S}}) \otimes \mathbb{Q} = N(\mathcal{S})$, where we write $\chi(T)$ for the character group of any torus T . Given any subtorus $U \subset A_{\mathcal{S}}$, we write $\mathrm{Ann}(U) \subset N(\mathcal{S})$ for the subspace of characters which annihilate U . This is a bijection between dimension d subtori of $A_{\mathcal{S}}$ and codimension d subspaces of $N(\mathcal{S})$. With this notation, $\mathrm{Ann}(T_{\mathcal{S}}) = \mathrm{Ann}(\Lambda^1)$.

Proposition 3.7. *For any subtorus $U \subset A_{\mathcal{S}}$, the intersection of U with the RM-torus $T_{\mathcal{S}}$ has codimension in $T_{\mathcal{S}}$ equal to $\dim \mathrm{ev}(\mathrm{Ann}(U))$, that is*

$$\dim(T_{\mathcal{S}}) - \dim(U \cap T_{\mathcal{S}}) = \dim \mathrm{ev}(\mathrm{Ann}(U)). \quad (3.2)$$

In particular,

$$\dim(T_{\mathcal{S}}) = \dim \mathrm{ev}(N(\mathcal{S})).$$

Proof. By Corollary 3.6, we have

$$\begin{aligned} \dim \mathrm{ev}(\mathrm{Ann}(U)) &= \dim \mathrm{Ann}(U) - \dim(\mathrm{Ann}(U) \cap \mathrm{Ann}(T_{\mathcal{S}})) \\ &= \dim(\mathrm{Ann}(U) + \mathrm{Ann}(T_{\mathcal{S}})) - \dim \mathrm{Ann}(T_{\mathcal{S}}). \end{aligned}$$

Also note that $\mathrm{Ann}(U \cap T_{\mathcal{S}}) = \mathrm{Ann}(U) + \mathrm{Ann}(T_{\mathcal{S}})$. It follows that

$$\begin{aligned} \dim \mathrm{ev}(\mathrm{Ann}(U)) &= \dim \mathrm{Ann}(U \cap T_{\mathcal{S}}) - \dim \mathrm{Ann}(T_{\mathcal{S}}) \\ &= \dim(T_{\mathcal{S}}) - \dim(U \cap T_{\mathcal{S}}). \end{aligned}$$

To obtain the last statement, apply (3.2) for U the trivial torus. ■

As an application, we can now calculate the dimension of $T_{\mathcal{S}}$ for many strata \mathcal{S} .

Proposition 3.8. *Suppose that \mathcal{S} is an admissible weighted boundary stratum for which the dual graph $\Gamma(\mathcal{S})$ contains an edge joining a vertex to itself. Then $\dim(T_{\mathcal{S}}) = g - 1$.*

Proof. By Propositions 3.4 and 3.7 we need only to show that $\dim \text{ev}(N(\mathcal{S})) \geq g - 1$. Let γ_1 be a loop which joins a vertex of $\Gamma(\mathcal{S})$ to itself, and let $K \subset F$ be the span of $\lambda(\gamma)$ over all loops γ which are edge-disjoint from γ_1 . Since λ induces an isomorphism $H_1(\Gamma(\mathcal{S}); \mathbb{Q}) \rightarrow F$, and deleting a loop from a graph reduces the rank of its homology by one, the dimension of K is $g - 1$. By Lemma 2.2, $\lambda(\gamma_1) \otimes K \in N(\mathcal{S})$, so $\lambda(\gamma_1)K \subset \text{ev}(N(\mathcal{S}))$. ■

4 Reduction to the boundary condition

In this section we show how the main theorem reduces to a containment statement about tori in the locus of stable forms for some boundary strata of $\overline{\mathcal{M}}_4$. For this purpose we call a boundary stratum \mathcal{S} of \mathcal{M}_4 *relevant*, if it parametrizes curves of geometric genus zero and if $\dim(\mathcal{S}) \geq 3$. Since for geometric genus zero curves the dimension of \mathcal{S} equals six minus the number of irreducible components of any stable curve of the stratum, the last condition is equivalent to having at most three irreducible components.

Proof of Theorem 1.1. Suppose that, contrary to the claim of the theorem, for some order \mathcal{O} and some symplectic \mathcal{O} -module M the component $\mathcal{R}\mathcal{A}_{\mathcal{O},M}$ of the real multiplication locus $\mathcal{R}\mathcal{A}_{\mathcal{O}}$ is generically contained in \mathcal{M}_4 . We denote by $\mathcal{R}\mathcal{M}_{\mathcal{O},M}$ the preimage of this component under the Torelli map. Since the Hilbert modular variety is not compact, the intersection $\partial\mathcal{R}\mathcal{M}_{\mathcal{O},M}$ of $\overline{\mathcal{R}\mathcal{M}_{\mathcal{O},M}}$ and the boundary part of the boundary of $\overline{\mathcal{M}}_g$ consisting of curves with non-compact Jacobian inside $\overline{\mathcal{M}}_g$ is non-empty. In fact $\partial\mathcal{R}\mathcal{M}_{\mathcal{O},M}$ must be a divisor on $\overline{\mathcal{R}\mathcal{M}_{\mathcal{O},M}}$, hence all irreducible components of $\partial\mathcal{R}\mathcal{M}_{\mathcal{O},M}$ are of dimension three. By [BM, Corollary 5.6], $\partial\mathcal{R}\mathcal{M}_{\mathcal{O},M}$ lies in the union of boundary strata parameterizing curves of geometric genus zero. More precisely, by Theorem 2.1 the boundary of $\mathcal{R}\mathcal{M}_{\mathcal{O},M}$ lies in the image of the $\mathcal{R}\mathcal{S}_E$ for some extension classes E .

All together, each irreducible component of $\partial\mathcal{R}\mathcal{M}_{\mathcal{O},M}$ generically lies in some relevant admissible weighted boundary stratum and for each the relevant admissible weighted boundary stratum \mathcal{S} that $\partial\mathcal{R}\mathcal{M}_{\mathcal{O},M}$ intersects, there is some extension class E such that $\mathcal{R}\mathcal{S}_E$ is of dimension three.

The following Propositions 4.1, 4.2 and 4.3 provide the contradiction we need to prove Theorem 1.1 for all the topological types of \mathcal{S} . ■

Proposition 4.1. *For each relevant weighted boundary stratum \mathcal{S} of $\overline{\mathcal{M}}_4$ without separating nodes, the topological type of \mathcal{S} being listed in Figure 1, the cross-ratio map CR is finite. In particular for each extension class E ,*

$$\dim \mathcal{R}\mathcal{S}_E = \dim(\text{CR}(\mathcal{S}) \cap T_{\mathcal{S},E}),$$

where the intersection is taken inside the ambient torus A_S .

Proposition 4.2. *For each relevant weighted boundary stratum \mathcal{S} of $\overline{\mathcal{M}}_4$ without separating nodes, the topological type of \mathcal{S} being listed in Figure 1, the intersection of $\text{CR}(\mathcal{S})$ with each translated cross-ratio torus $T_{S,E}$ inside the ambient torus is of dimension at most two.*

It remains to show that a component of $\mathcal{RA}_{\mathcal{O}}$ contained in \mathcal{M}_4 cannot only meet boundary strata which have separating curves.

Proposition 4.3. *Suppose that a component $\mathcal{RA}_{\mathcal{O},M}$ of the real multiplication locus is generically contained in the Torelli image of $\overline{\mathcal{M}}_4$. Suppose moreover, that the closure of $\mathcal{RM}_{\mathcal{O},M}$ intersects the image in $\overline{\mathcal{M}}_4$ of a weighted relevant boundary stratum \mathcal{S} parameterizing stable curves with a separating node. Then there exists also an irreducible component of $\partial\mathcal{RM}_{\mathcal{O},M}$ contained in a relevant boundary stratum parameterizing stable curves without a separating node.*

Proposition 4.1 is a weaker version of Theorem 5.1. The proof of the other two propositions will occupy the rest of the paper.

Relevant boundary strata of \mathcal{M}_4 . Figure 1 contains the complete list of relevant boundary strata of \mathcal{M}_4 parameterizing stable curves without a separating node. We will refer to the stratum (x, y) as the stratum in row x and column y . The strata $(2, 2)$ (“the $[5] \times^5 [5]$ -stratum”) and $(4, 2)$ (“the doubled triangle”) will need a special treatment below.

The arrows are chosen arbitrarily. Their purpose is to label the marked points on the normalization of the stable curve that are glued together. Our convention is that on the edge with label k the points P_k and Q_k are glued together, where the point P_k sits on the outgoing component and Q_k sits on the incoming component. If the graph is given an F -weighting, we call r_k the weight of the k th edge.

There are many choices for labeling of the edges. Our choices have the property that the first four weights always span F :

Lemma 4.4. *Consider an F -weighted graph Γ containing edges e_1, \dots, e_4 having weights w_1, \dots, w_4 . Then the weights w_i span F if and only if the complement of the e_i in Γ is a tree.*

In particular, with the choice of labeling given in Figure 1, every relevant F -weighted boundary stratum with no separating nodes has the property that the weights $\{r_1, r_2, r_3, r_4\}$ are a \mathbb{Q} -basis of F .

Proof. Recall that there is the isomorphism $\lambda: H_1(\Gamma; \mathbb{Q}) \rightarrow F$. Let $A \subset F$ be the annihilator of the span of the w_i with respect to the trace pairing. If the complement of the e_i contains a loop γ , then by the definition of λ , the nonzero element $\lambda(\gamma)$ pairs trivially with each w_i . Thus A is nontrivial, and the w_i do not span. The converse follows similarly. ■

Consequently, we may set (s_1, s_2, s_3, s_4) to be the basis of F dual to the basis (r_1, r_2, r_3, r_4) , and we keep this notation throughout the rest of the paper.




Figure 1: Relevant genus four stable curves without separating nodes

5 Cross-ratios for nice boundary strata without separating nodes

The next theorem is completely analogous to [BM, Proposition 8.3], except that we now work in genus 4 instead of genus 3. It is a simple Torelli type theorem for relevant boundary strata without separating nodes.

For any weighted boundary stratum \mathcal{S} let ϕ_ι be the involution which changes each weight to its negative. We say that \mathcal{S} is ϕ_ι -invariant, if $\phi_\iota(\mathcal{S})$ and \mathcal{S} are topologically equivalent, in which case ϕ_ι restricts to an involution of \mathcal{S} . Among the relevant boundary strata without separating nodes, the ϕ_ι -invariant strata are precisely the strata (1, 1), (1, 2), (2, 2), (2, 3), and (3, 3). Note that whether or not a weighted stratum is ϕ_ι -invariant depends only on its topological type and not on the choice of weights.

By the definition of ϕ_ι , each period matrix entry $\Psi(x)$ is equivariant with respect to the involution ϕ_ι . If \mathcal{S} is ϕ_ι -invariant, we define \mathcal{S}' to be the quotient of \mathcal{S} by ϕ_ι .

Recall from §2 that we have a canonical morphism CR: $\mathcal{S} \rightarrow A_{\mathcal{S}}$. A basis τ_1, \dots, τ_n of $N(\mathcal{S})$ determines an isomorphism of $A_{\mathcal{S}}$ with $(\mathbb{C}^*)^n$, and in these coordinates, CR is simply the product of the functions $\Psi(\tau_i)$. By the above discussion, CR factors through ϕ_ι to define a morphism CR: $\mathcal{S}' \rightarrow A_{\mathcal{S}}$.

Theorem 5.1. *Given a relevant stratum without separating nodes \mathcal{S} , the morphism CR is an embedding of \mathcal{S} (or \mathcal{S}' if \mathcal{S} is ϕ_ι -invariant) in $A_{\mathcal{S}}$.*

Proof. We give the details for some strata where essential arguments show up and leave the remaining verifications to the reader.

We first consider the irreducible stratum (1, 1). This stratum is ϕ_ι -invariant, and the involution has the effect of swapping each pair P_i, Q_i . Here $s_i \otimes s_j \in N(\mathcal{S})$ for all $i \neq j$ and we abbreviate $\Psi(s_i \otimes s_j) = R_{ij}$ where

$$R_{ij} = [P_i, Q_i, P_j, Q_j]$$

by Lemma 2.2 or [BM, Proposition 8.3]. We fix $P_1 = 0, Q_1 = \infty$ and $P_2 = 1$. We then have $R_{12} = Q_2$ and $P_3 R_{13} = Q_3$. Given R_{23} , we need to solve a quadratic equation to recover P_3 and Q_3 . Similarly, given R_{14} and R_{24} we need to solve a quadratic equation to recover P_4 and Q_4 . Thus there are four possibilities for the tuple (P_3, Q_3, P_4, Q_4) . The cross-ratio R_{34} eliminates two of these solutions, and the remaining two are related by the involution ϕ_ι .

Next we consider the “[5] \times^5 [5]”-stratum (2, 2). Again, ϕ_ι swaps each pair P_i, Q_i . Here $s_i \otimes (s_j - s_k) \in N(\mathcal{S})$ for all distinct i, j, k . We normalize

$$P_1 = Q_1 = 1, \quad P_2 = Q_2 = 0, \quad P_3 = Q_3 = \infty.$$

Then

$$\begin{aligned} \Psi(s_1 \otimes (s_4 - s_2)) &= (1 - P_4)(1 - Q_4), & \Psi(s_2 \otimes (s_4 - s_1)) &= P_4 Q_4 \\ \Psi(s_1 \otimes (s_3 - s_2)) &= (1 - P_3)(1 - Q_3), & \Psi(s_2 \otimes (s_3 - s_1)) &= P_3 Q_3. \end{aligned}$$

The first line determines two possibilities for P_4 and Q_4 , and the second line determines two possibilities for P_3 and Q_3 . Using $\Psi(s_3 \otimes (s_1 - s_2))$ eliminates two of the four possibilities for (P_3, P_4, Q_3, Q_4) , with the remaining two possibilities related by ϕ_ι .

As typical examples for the remaining cases we take the stratum $(1, 4)$. Here $N(\mathcal{S})$ is generated by $s_1 \otimes s_2$, $s_1 \otimes s_3$, $s_4 \otimes s_2$ and $s_4 \otimes s_3$. Normalizing

$$P_1 = P_4 = 0, \quad Q_1 = Q_4 = \infty, \quad P_2 = Q_2 = 1$$

we obtain

$$\begin{aligned} \Psi(-s_1 \otimes s_2) &= Q_5, & \Psi(-s_1 \otimes s_3) &= Q_3, \\ \Psi(-s_4 \otimes s_2) &= P_5, & \Psi(-s_4 \otimes s_3) &= P_3, \end{aligned}$$

so these four cross-ratios determine the remaining four points. ■

Gerritzen's equation. Given the above Torelli theorem, it is natural to ask what the image of CR in the ambient torus is. For the irreducible stratum $(1, 1)$ the question of finding the equation cutting out the image of CR has been solved by [Ger92]. For all but one exceptional stratum, we will be able to avoid the use of this equation. For one exceptional stratum we indeed need to determine the image of CR and this equation can be obtained as a limit of Gerritzen's equation.

Proposition 5.2 ([Ger92, Proposition 4.3.1]). *For the irreducible stratum, the image of CR in the ambient torus $\mathcal{A}_{\mathcal{S}}$ is given, in the coordinates introduced in the proof of Theorem 5.1, as the vanishing locus of the function $F = \Delta H - G$, where*

$$\begin{aligned} \Delta &= (R_{12} - 1)(R_{13} - 1)(R_{14} - 1)(R_{23} - 1)(R_{24} - 1)(R_{34} - 1) \\ H &= R_{12}R_{13}R_{14}R_{23}R_{24}R_{34} - R_{12}R_{14}R_{24} - R_{13}R_{14}R_{34} - R_{23}R_{24}R_{34} \\ &\quad - R_{12}R_{13}R_{23} + R_{14}R_{23} + R_{13}R_{24} + R_{12}R_{34} \\ G &= R_{12}R_{34}(R_{13} - 1)^2(R_{14} - 1)^2(R_{23} - 1)^2(R_{24} - 1)^2 \\ &\quad + R_{13}R_{24}(R_{12} - 1)^2(R_{14} - 1)^2(R_{23} - 1)^2(R_{34} - 1)^2 \\ &\quad + R_{14}R_{23}(R_{12} - 1)^2(R_{13} - 1)^2(R_{24} - 1)^2(R_{34} - 1)^2. \end{aligned}$$

The validity of the equation, hence the fact that we use the same conventions on cross-ratios as Gerritzen, can be checked by plugging in the definition of the cross-ratios.

A hypothetical sufficiency theorem. In [BM], we proved that the necessary condition of Theorem 2.1 is also sufficient in genus three. The proof relied heavily on the fact that the Schottky problem is trivial in genus three, that is, the image of the Torelli map $\mathcal{M}_g \rightarrow \mathcal{A}_g$ is dense. In higher genus, we do not know whether our condition is also sufficient; however, under the assumption that a component $\mathcal{R}\mathcal{A}_{\mathcal{O}, M}$ of the real multiplication is contained in the Schottky locus sufficiency holds at least in some cases.

We say that a stable curve is *nice* if the complement of any two nodes is connected (such curves are sometimes called three-connected). A boundary stratum is nice if it consists of nice stable curves.

Theorem 5.3. *Assuming that a component $\mathcal{RA}_{\mathcal{O},M}$ is contained in the closure of $t(\mathcal{M}_4)$, the necessary condition of Theorem 2.1 for geometric genus zero stable curve to lie in the boundary of $\mathcal{RM}_{\mathcal{O},M}$ is also sufficient for nice stable curves.*

We emphasize that this theorem has no use outside of this paper, as we are proving that the hypothesis of the theorem never holds. As the proof of Theorem 5.3 is essentially the same as in [BM], we only sketch the idea of the proof here.

Consider a nice boundary stratum \mathcal{S} and choose a basis τ_1, \dots, τ_n of $N(\mathcal{S})$. By Theorem 5.1, the map $\text{CR}: \mathcal{S} \rightarrow (\mathbb{C}^*)^n$ is either two-to-one or biholomorphic onto its image $\text{CR}(\mathcal{S})$. In either case this map is open. In genus three, this map is open as well and is also onto. In [BM], we extended this to an open map $\Xi: U \rightarrow \mathbb{C}^m \times (\mathbb{C}^*)^n$ (for some neighborhood U of \mathcal{S}) sending \mathcal{S} to $\{\mathbf{0}\} \times (\mathbb{C}^*)^n$. In genus four, the map Ξ is defined in the same way and is also an open map $\Xi: U \rightarrow \mathbb{C}^m \times \text{CR}(\mathcal{S}) \subset \mathbb{C}^m \times (\mathbb{C}^*)^n$. In order to define the map Ξ , it is necessary for \mathcal{S} to be nice.

The map Ξ sends a component of the real multiplication locus to a subvariety $T \subset (\mathbb{C}^*)^m \times (\mathbb{C}^*)^n$ which is a translate of a torus. As Ξ is open, to show that a point $p \in \mathcal{S}$ is in the boundary of the real multiplication locus, it suffices to show that $\Xi(p)$ is in the boundary of T . Thus the problem is reduced to calculating the boundary of a torus. In [BM, Theorem 8.14], we construct an explicit one-dimensional torus $T_1 \subset T$ which limits on $\Xi(p)$, showing that $\Xi(p)$ is indeed in the boundary of T .

In genus three, this proof uses in an essential way that a generic abelian variety is a Jacobian. In higher genus, this proof breaks down, since we don't know whether the torus T_1 is contained in the Schottky locus. However, assuming that the component of the real multiplication we are considering is contained in the Schottky locus, this is automatic and the proof carries through.

6 Checking nice boundary strata without separating nodes

In this section we develop two criteria on the dual graphs of strata to test whether Proposition 4.2 holds. These criteria will apply to each of the strata in Figure 1 except for two exceptional strata which we handle with *ad hoc* arguments.

In the following, it will be useful to encode loops on such a dual graph by the labeling of the edges. We use the convention that the first digit corresponds to the edge used first and an overline corresponds to using the edge in the direction pointing opposite the arrow. E.g. in the stratum (1, 2), the loop (3 $\overline{5}$) turns the counterclockwise around the middle circle, starting at the upper vertex.

The disjoint loop argument. The first criterion rules out strata whose dual graphs contain disjoint loops:

Proposition 6.1. *Let $\Gamma(\mathcal{S})$ be the dual graph of a relevant nice boundary stratum without separating nodes \mathcal{S} . Suppose Γ contains two vertex-disjoint simple loops γ_1 and γ_2 . Then Proposition 4.2 holds for \mathcal{S} .*

Proof. Suppose the contrary holds, i.e. $T_{\mathcal{S},E} \subset \overline{\text{CR}(\mathcal{S})}$. Let $a = \lambda(\gamma_1) \otimes \lambda(\gamma_2) \in N(\mathcal{S})$ as in Lemma 2.2. We claim that $\text{ev}(a) \neq 0$. To justify this, it suffices to show that the field element associated with any simple loop is non-zero. This is a consequence of the isomorphism $\lambda : H_1(\Gamma(\mathcal{S}), \mathbb{Z}) \rightarrow \mathcal{I}^\vee$ and the fact that such a loop is non-zero in $H_1(\Gamma(\mathcal{S}), \mathbb{Z})$.

Since the loops are vertex-disjoint, the Lemma 2.2 implies $\Psi(a) \equiv 1$, i.e. $T_{\mathcal{S},E}$ is contained in the torus U with $\text{Ann}(U) = \langle a \rangle$. Together with $\text{ev}(a) \neq 0$ this contradicts Proposition 3.7. ■

The shared vertex argument. Recall that a graph is called n -connected if it can not be disconnected by removing $n - 1$ edges. We say that it is *precisely n -connected* if it is n -connected and can be disconnected by removing n edges. Note that the disjoint loop argument applies to any precisely 2-connected stratum.

Proposition 6.2. *Proposition 4.2 holds for a relevant nice boundary stratum without separating nodes \mathcal{S} provided the following condition on the dual graph $\Gamma(\mathcal{S})$ holds.*

Suppose $\Gamma(\mathcal{S})$ contains two edge-disjoint loops γ_1 and γ_2 having exactly one vertex v in common. Suppose moreover, that there is some precisely 2-connected graph Γ' obtained from $\Gamma(\mathcal{S})$ by gluing an edge e into v such that γ_1 and γ_2 yield vertex-disjoint loops in Γ' , and moreover for all such graphs Γ' the following condition holds. There is a loop γ_3 on Γ' such that γ_3 and γ_1 or γ_3 and γ_2 are vertex-disjoint.

Proof. We let $\tilde{\mathcal{S}}$ be the partial Deligne-Mumford compactification of \mathcal{S} obtained by adjoining any weighted stable curve which has the same set of weights as the curves parametrized by \mathcal{S} . For each degeneration \mathcal{S}' in $\tilde{\mathcal{S}} \setminus \mathcal{S}$, we then have $N(\mathcal{S}') = N(\mathcal{S})$, so the morphism CR extends to a morphism $\text{CR} : \tilde{\mathcal{S}} \rightarrow A_{\mathcal{S}}$.

Now suppose the contrary holds, i.e. $T_{\mathcal{S},E} \subset \overline{\text{CR}(\mathcal{S})}$. Let $a = \lambda(\gamma_1) \otimes \lambda(\gamma_2) \in N(\mathcal{S})$ and consider the intersection of $\text{CR}(\tilde{\mathcal{S}})$ with the subtorus U given by $\text{Ann}(U) = \langle a \rangle$. This intersection is nonempty and consists of the union of all degenerations $\text{CR}(\mathcal{S}')$ which correspond to some graph Γ' as in the statement. By Proposition 3.7, the intersection $U \cap T_{\mathcal{S},E}$ is a translate of a two-dimensional subtorus of $A_{\mathcal{S}}$, thus it is contained in one degeneration $\text{CR}(\mathcal{S}')$ of $\text{CR}(\mathcal{S})$ as above. In what follows we fix this degeneration \mathcal{S}' .

Let $b = \lambda(\gamma_1) \otimes \lambda(\gamma_3)$ resp. $\lambda(\gamma_3) \otimes \lambda(\gamma_2)$ depending on which loops are disjoint on \mathcal{S}' . The preceding argument together with Lemma 2.2 imply that on $\text{CR}(\mathcal{S}') \cap U$ the function $\Psi(b)$ is identically one. Consider the torus U_2 defined by $\text{Ann}(U_2) = \langle a, b \rangle$. Since $\Psi(b) \equiv 1$ on $\text{CR}(\mathcal{S}')$, we have $U_2 \cap T_{\mathcal{S},E} =$

$U \cap T_{\mathcal{S},E}$, so this intersection is two-dimensional. Since $\dim \text{ev}(\text{Ann}(U_2)) = 2$, this contradicts Proposition 3.7. \blacksquare

These two arguments allow us to prove Proposition 4.2 in all but two cases. The disjoint loop argument applies to the 2-connected strata $(1, 2)$, $(1, 4)$, $(2, 3)$, $(2, 4)$, $(3, 1)$, $(3, 2)$, $(3, 3)$, $(4, 1)$, and $(4, 3)$.

To deal with the stratum $(1, 1)$ we apply the shared vertex argument to $\gamma_1 = (1)$ and $\gamma_2 = (2)$. There is only one precisely 2-connected degeneration, namely the stratum $(1, 2)$. Obviously γ_3 with the required properties exists.

To deal with the stratum $(1, 3)$ we apply the shared vertex argument to $\gamma_1 = (1)$ and $\gamma_2 = (3\bar{4})$. There are three precisely 2-connected degenerations that make γ_1 and γ_2 disjoint. The reader will check easily that in all three cases a loop γ_3 with the required properties exists.

To deal with the stratum $(2, 1)$ we use the loops $\gamma_1 = (1)$ and $\gamma_2 = (2\bar{3})$ for the shared vertex argument. To deal with the stratum $(2, 5)$ we use the loops $\gamma_1 = (1\bar{2})$ and $\gamma_2 = (3\bar{4})$. To deal with the stratum $(3, 4)$ we use the loops $\gamma_1 = (3\bar{6})$ and $\gamma_2 = (2\bar{5})$ for the shared vertex argument. To deal with the stratum $(4, 4)$ we use the loops $\gamma_1 = (1\bar{5})$ and $\gamma_2 = (2\bar{6})$. In all these cases, there is only one precisely 2-connected degeneration, and the required γ_3 exists.

The exceptional cases 'doubled triangle' and $[5] \times^5 [5]$. Finally, we treat two exceptional cases separately.

Proposition 6.3. *Proposition 4.2 holds for the " $[5] \times^5 [5]$ -stratum" given by the graph $(2, 2)$.*

Proof. A basis of $N(\mathcal{S})$ for this stratum is given by $s_i \otimes s_j - s_3 \otimes s_4$, with $i < j$ and $(i, j) \neq (3, 4)$. We view this stratum as a degeneration of the irreducible stratum, obtained by unpinching the node with label 5. We derive the equation of the CR-image of this stratum from the equation of the irreducible stratum in Proposition 5.2. The coordinates $\tilde{R}_{ij} = \Psi(s_i \otimes s_j - s_3 \otimes s_4)$ are related to the coordinates in that proposition by $\tilde{R}_{ij} = R_{ij}/R_{34}$. Pinching the node with label 5 takes R_{34} to ∞ . Hence in order to determine the image of CR for the $[5] \times^5 [5]$ -stratum, we rewrite Gerritzen's equation in terms of the \tilde{R}_{ij} and R_{34} , and consider the leading term for $R_{34} \rightarrow \infty$.

Consequently, in these coordinates the image of CR is the variety $V(F_5)$ cut out by $F_5 = 0$, where

$$F_5 = \tilde{R}_{12}\tilde{R}_{13}\tilde{R}_{14}\tilde{R}_{23}\tilde{R}_{24} - \tilde{R}_{12}\tilde{R}_{14}\tilde{R}_{23} - \tilde{R}_{12}\tilde{R}_{13}\tilde{R}_{24} - \tilde{R}_{13}\tilde{R}_{14}\tilde{R}_{23}\tilde{R}_{24}.$$

For convenience, we relabel the coordinates \tilde{R}_{ij} as Z_1, \dots, Z_5 , using the lexicographical order.

Consider the vectors of exponents $v_1 = (1, 1, 1, 1, 1)$, $v_2 = (1, 0, 1, 1, 0)$, $v_3 = (1, 1, 0, 0, 1)$, $v_4 = (0, 1, 1, 1, 1)$ for the monomials m_1, \dots, m_4 appearing in F_5 . Suppose first that $V(F_5)$ contains a translate of the torus T parametrized by $f_{\mathbf{a}}(t) = (t^{a_1}, \dots, t^{a_5})$. We have $m_i \circ f(t) = t^{n_i}$, where $n_i = v_i \cdot \mathbf{a}$. It follows that

for T to be contained in $V(F_5)$, we must have that for each v_i , there is some other v_j such that $(v_i - v_j) \cdot \mathbf{a} = 0$.

Now suppose F_5 contains a translate of the three-dimensional torus $T_{\mathcal{S}}$. Let $P \subset \mathbb{Q}^5$ be the three-dimensional subspace which parametrizes T , and let $N = P^\perp \subset \mathbb{Q}^5$. Given v_i , by the above discussion there must be some $j \neq i$ such that $v_i - v_j \in N$ (using that a vector space over \mathbb{Q} can not be the union of proper subspaces). Since the span of $\{v_1, v_2, v_3, v_4\}$ is three-dimensional, this is only possible if there is a basis n_1, n_2 of N such that n_1 is the difference of two of the v_i and n_2 is the difference of the other two v_i . Suppose that $n_1 = v_2 - v_1$ and $n_2 = v_3 - v_4$, the other two cases will lead to the same contradiction.

By Proposition 3.5 the condition that n_2 is perpendicular to $T_{\mathcal{S}}$ is equivalent to

$$\text{Tr}(x(s_1s_2 - s_1s_4 - s_2s_3 + s_3s_4)) = 0 \quad \text{for all } x \in F,$$

i.e. $0 = s_1s_2 - s_1s_4 - s_2s_3 + s_3s_4 = (s_1 - s_3)(s_2 - s_4)$. This contradicts that the s_i are a \mathbb{Q} -basis of F . ■

Proposition 6.4. *Proposition 4.2 holds for the “doubled triangle-stratum” \mathcal{S} given by the graph (4, 2).*

Proof. By Lemma 2.2 the three pairs of loops $((1\bar{4}), (3\bar{6}))$, $((2\bar{5}), (1\bar{4}))$ and $((3\bar{6}), (2\bar{5}))$ define elements of $N(\mathcal{S})$. Their Ψ -images are

$$R_1 = [P_1, Q_3, Q_6, P_4], \quad R_2 = [P_2, Q_1, Q_4, P_5], \quad R_3 = [P_3, Q_2, Q_5, P_6].$$

Consider now the loops (123) and (456). By the same lemma they define an element in $N(\mathcal{S})$ whose Ψ -image is

$$R_4 = [P_1, Q_6, P_4, Q_3][P_2, Q_4, P_5, Q_1][P_3, Q_5, P_6, Q_2] = \frac{1}{1 - R_1} \frac{1}{1 - R_2} \frac{1}{1 - R_3}$$

Since \mathcal{S} is irreducible and $\dim \text{CR}(\mathcal{S}) = 3$, if $\text{CR}(\mathcal{S})$ contains the RM-torus, then $\text{CR}(\mathcal{S})$ is equal to that torus. But the above equation obviously does not cut out a subtorus of the ambient torus $A_{\mathcal{S}}$ with coordinates R_1, R_2, R_3, R_4 . ■

7 Strata with separating curves

Finally, we prove Proposition 4.3, completing the proof of Theorem 1.1.

Proof of Proposition 4.3. By Theorem 2.1 there is an admissible, \mathcal{I} -weighted boundary stratum \mathcal{S}' such that the boundary component of $\mathcal{RM}_{\mathcal{O}, M}$ given by hypothesis lies in the image of \mathcal{S}' under the forgetful map $\mathcal{S}' \rightarrow \overline{\mathcal{M}}_4$. Moreover, by hypothesis, the dual weighted graph $\Gamma' = \Gamma(\mathcal{S}')$ has a separating edge e . Let $\tilde{\Gamma}$ be the weighted graph obtained by contracting the separating edges of Γ' , preserving weights on the other edges. If $\tilde{\Gamma}$ happens to be not nice, say the pair of edges f and g disconnects $\tilde{\Gamma}$, then we contract g preserving weights on the other edges. The resulting weighted graph will still be admissible, since the

weights on f and g are r and $-r$. We keep contracting edges until we arrive at a nice graph Γ . If $\tilde{\Gamma}$ was nice to start with, we take $\Gamma = \tilde{\Gamma}$.

Let \mathcal{S} be the corresponding boundary stratum and let $\bar{\mathcal{S}}$ be the partial Deligne-Mumford compactification, adding those stable curves whose dual graphs lie between Γ' and Γ . We want to intersect the cross-ratio images of these spaces with the translated RM-torus $T_{\mathcal{S},E}$. The situation is summarized in the following diagram, where we emphasize that the cross-ratio map on left is not injective but the one in the middle is injective or two-to-one.

$$\begin{array}{ccccc}
 \mathcal{S}' & \hookrightarrow & \bar{\mathcal{S}} & & T_{\mathcal{S},E} \\
 \downarrow \text{CR} & & \downarrow \text{CR} & \searrow \text{CR} & \downarrow \\
 \text{CR}(\mathcal{S}') & \hookrightarrow & \text{CR}(\bar{\mathcal{S}}) & \hookrightarrow & A_{\mathcal{S}}
 \end{array}$$

We may restrict ourselves the case $\dim \mathcal{S} \geq 4$, since otherwise \mathcal{S}' cannot contain a boundary divisor of the real multiplication locus. For the stratum $[5] \times^5 [5]$ there is no codimension-one degeneration with a separating edge. Since all the other strata \mathcal{S} with $\dim \mathcal{S} \geq 4$ have a loop joining a node to itself, we conclude from Proposition 3.8 that $\dim T_{\mathcal{S},E} = 3$.

The separating edge may split the stable curves into two components either of genera 2 and 2 (the (2,2)-case) or of genera 1 and 3 (the (1,3)-case). A separating edge also defines a splitting of F into two \mathbb{Q} -subspaces F_1 and F_2 generated by the λ -images of loops in the components of $\Gamma' \setminus \{e\}$. Each element $a \in F_1 \otimes F_2$ defines by Lemma 2.2 an element of $N(\mathcal{S})$, and $\text{CR}(\mathcal{S}')$ is contained in the subtorus defined by $\psi(a) = 1$ for all $a \in F_1 \otimes F_2$.

We claim that it is enough to show that $T_{\mathcal{S},E} \cap \text{CR}(\mathcal{S}) \neq \emptyset$. Suppose that this intersection is in fact nonempty. By the sufficiency criterion Theorem 5.3, this intersection belongs to the intersection of $\overline{\mathcal{RM}}_{\mathcal{O},M}$ with the boundary of $\overline{\mathcal{M}}_4$. As $\dim \mathcal{RM}_{\mathcal{O},M} = 4$, each irreducible component of this intersection must be three dimensional. Thus $T_{\mathcal{S},E} \cap \text{CR}(\mathcal{S})$ is contained in a three-dimensional component of $\partial \mathcal{RM}_{\mathcal{O},M}$ which lies in some stratum (possibly obtained by further undegenerating \mathcal{S}) without separating nodes.

We start with the case of the irreducible stratum \mathcal{S} , hence $\dim A_{\mathcal{S}} = 6$. By the above discussion, we must show that the intersection $T_{\mathcal{S},E} \cap \text{CR}(\bar{\mathcal{S}})$ is not contained in $\text{CR}(\mathcal{S}')$. This intersection is at least two-dimensional, so it suffices to show that $T_{\mathcal{S},E} \cap \text{CR}(\mathcal{S}')$ is at most one-dimensional. In the (2,2)-case $\dim(F_1 \otimes F_2) = 4$, hence $\text{CR}(\mathcal{S}') = \text{CR}(\mathcal{S}') \cap T_{\mathcal{S},E}$. Proposition 3.7 applied to the torus $U = \text{CR}(\mathcal{S}')$ and $\dim \text{ev}(F_1 \otimes F_2) \geq 2$ shows that this intersection is one-dimensional. In the (1,3)-case $\dim(F_1 \otimes F_2) = 3$ and $\dim \text{CR}(\mathcal{S}')$ is at least three by the genus three analog of Theorem 5.1 [BM, Corollary 8.4], hence $\text{CR}(\mathcal{S}')$ coincides with the subtorus of $A_{\mathcal{S}}$ cut out by $F_1 \otimes F_2 \subset N(\mathcal{S})$. We now apply Proposition 3.7 to the torus $U = \text{CR}(\mathcal{S}')$ to show that the intersection with $T_{\mathcal{S},E}$ is one-dimensional.

If the stratum \mathcal{S} is reducible, we have $\dim \mathcal{S} = 4$ and $\dim A_{\mathcal{S}} = 5$. Again we must show that $T_{\mathcal{S},E} \cap \text{CR}(\mathcal{S}')$ is at most one-dimensional. In the (2,2)-case $\dim(F_1 \otimes F_2) = 4$, hence the codimension of $\text{CR}(\mathcal{S}')$ in the ambient torus

A_S is at least 4 and we obtain immediately a contradiction. In the $(1, 3)$ -case $\dim(F_1 \otimes F_2) = 3$, hence $\text{CR}(S')$ has to be a 2-dimensional torus to which we apply again Proposition 3.7. Since $\dim \text{ev}(F_1 \otimes F_2) = 3 > 1$, we again have a contradiction. ■

References

- [Ale04] Valery Alexeev. Compactified Jacobians and Torelli map. *Publ. Res. Inst. Math. Sci.*, 40(4):1241–1265, 2004.
- [Bai07] M. Bainbridge. Euler characteristics of Teichmüller curves in genus two. *Geom. Topol.*, 11:1887–2073, 2007.
- [BM] M. Bainbridge and M. Möller. The Deligne-Mumford compactification of the real multiplication locus and Teichmüller curves in genus three, arXiv:math.AG/0911.4677, to appear in *Acta Math.*
- [BM76] J. Bondy and U. Murty. *Graph theory with applications*. American Elsevier Publishing Co., Inc., New York, 1976.
- [dJZ07] J. de Jong and S. Zhang. Generic abelian varieties with real multiplication are not Jacobians. In *Diophantine geometry*, volume 4 of *CRM Series*, pages 165–172. Ed. Norm., Pisa, 2007.
- [FM98] B. Farb and H. Masur. Superrigidity and mapping class groups. *Topology*, 37(6):1169–1176, 1998.
- [Ger92] L. Gerritzen. Equations defining the periods of totally degenerate curves. *Israel J. Math.*, 77(1-2):187–210, 1992.
- [Hai99] R. Hain. Locally symmetric families of curves and Jacobians. In *Moduli of curves and abelian varieties*, *Aspects Math.*, E33, pages 91–108. Vieweg, Braunschweig, 1999.
- [MO] B. Moonen and F. Oort. The Torelli locus and special subvarieties The theta characteristic of a branched covering, Preprint, August 2010, 43pp. Submitted to the Handbook of Moduli (G. Farkas and I. Morrison, eds.).
- [MZ11] M. Möller and D. Zagier. Theta derivatives and Teichmüller curves, 2011, in preparation, abridged version on the the web page of M. Möller.
- [Nam76] Yukihiro Namikawa. A new compactification of the Siegel space and degeneration of Abelian varieties. I. *Math. Ann.*, 221(2):97–141, 1976.